

Hypertension

JOURNAL OF THE AMERICAN HEART ASSOCIATION



*Learn and Live*SM

Central Blood Pressure Measurements and Antihypertensive Therapy: A Consensus Document

Enrico Agabiti-Rosei, Giuseppe Mancia, Michael F. O'Rourke, Mary J. Roman, Michel E. Safar, Harold Smulyan, Ji-Guang Wang, Ian B. Wilkinson, Bryan Williams and Charalambos Vlachopoulos

Hypertension 2007;50;154-160; originally published online Jun 11, 2007;

DOI: 10.1161/HYPERTENSIONAHA.107.090068

Hypertension is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75214

Copyright © 2007 American Heart Association. All rights reserved. Print ISSN: 0194-911X. Online ISSN: 1524-4563

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://hyper.ahajournals.org/cgi/content/full/50/1/154>

Data Supplement (unedited) at:

<http://hyper.ahajournals.org/cgi/content/full/HYPERTENSIONAHA.107.090068/DC1>

Subscriptions: Information about subscribing to Hypertension is online at
<http://hyper.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, a division of Wolters Kluwer Health, 351 West Camden Street, Baltimore, MD 21202-2436. Phone: 410-528-4050. Fax: 410-528-8550. E-mail:
journalpermissions@lww.com

Reprints: Information about reprints can be found online at
<http://www.lww.com/reprints>

Central Blood Pressure Measurements and Antihypertensive Therapy

A Consensus Document

Enrico Agabiti-Rosei, Giuseppe Mancia, Michael F. O'Rourke, Mary J. Roman, Michel E. Safar, Harold Smulyan, Ji-Guang Wang, Ian B. Wilkinson, Bryan Williams, Charalambos Vlachopoulos

The 2003 European Society of Hypertension/European Society of Cardiology guidelines for the management of arterial hypertension¹ included 2 important novel recommendations: assessment of the total cardiovascular risk should be taken into account in the management of the hypertensive patient, and quantification of risk should include subclinical target organ damage.

These guidelines acknowledged that central (aortic) blood pressure (BP), which is the pressure exerted on the heart and brain, may be different from the pressure that is measured at the arm. They also recognized that central pressure may be predictive of outcome in specific populations² and differently affected by antihypertensive drugs. However, although these guidelines accepted that central augmentation index and pulse wave velocity may be important as measures of subclinical organ damage, they also stressed the need for prospective trials to establish their predictive values given that such studies were lacking at that time (2003).

After publication of these guidelines, additional data have strengthened the pathophysiological importance of central BP. Clinical studies have indicated that central BP may have predictive value independent of the corresponding peripheral (brachial) BP. More importantly, recent large-scale trials have shown that central hemodynamics may provide a worthwhile treatment target. In addition, central hemodynamics can now be reliably assessed noninvasively with a number of devices. Accordingly, because arterial stiffening and central hemodynamics are markers and manifestations of organ damage, the pertinent key question is whether the balance of evidence on their importance and issues related to clinical practice allows for implementation in patient management.

Pathophysiological Significance of Central Pressures

Central (aortic and carotid) pressures are pathophysiologically more relevant than peripheral pressures for the pathogenesis of cardiovascular disease.^{3,4} It is aortic systolic pressure that the left ventricle encounters during systole (afterload), and the aortic pressure during diastole is a determinant of coronary perfusion. Furthermore, the distending pressure in the large elastic-type arteries (aorta and carotid) is a key determinant of the degenerative changes that characterize accelerated aging and hypertension. In contrast, the muscular peripheral arteries, such as the brachial and the radial ones, are less influenced by these changes.⁵

The pressure wave generated by the left ventricle travels down the arterial tree and then is reflected at multiple peripheral sites, mainly at resistance arteries (small muscular arteries and arterioles). Consequently, the pressure waveform recorded at any site of the arterial tree is the sum of the forward traveling waveform generated by left ventricular ejection and the backward traveling wave, the "echo" of the incident wave reflected at peripheral sites. When the large conduit arteries are healthy and compliant, the reflected wave merges with the incident in the proximal aorta during diastole, thereby augmenting the diastolic BP and aiding coronary perfusion. In contrast, when the arteries are stiff, pulse wave velocity increases, accelerating the incident and reflected waves; thus, the reflected wave merges with the incident wave in systole and augments aortic systolic rather than diastolic pressure. As a result, left ventricular afterload increases, and normal ventricular relaxation and coronary filling are compromised. Apart from changes in the timing of the waveforms merging, changes in the magnitude of the reflected wave and central pressures may result from changes in the proportion of the incident wave that is reflected, which

Received February 27, 2007; first decision March 22, 2007; revision accepted April 25, 2007.

From the Clinica Medica and Department of Medicine (E.A.-R.), University Hospital, Brescia, Italy; Clinica Medica and Department of Medicine (G.M.), University of Milano-Bicocca, St Gerardo Hospital, Monza, Milan, Italy; St Vincent's Clinic/University of New South Wales (M.F.O.), Darlinghurst, Australia; Weill Medical College (M.J.R.), Cornell University, Ithaca, NY; Centre de Diagnostic (M.E.S.), Hopital Hotel-Dieu, Paris, France; Department of Medicine (H.S.), Upstate Medical University, Syracuse, NY; Centre for Epidemiological Studies and Clinical Trials (J.G.W.), Ruijin Hospital Shanghai Jiaotong, University School of Medicine, Shanghai, China; University of Cambridge Addenbrooke's Hospital (I.B.W.), Cambridge, United Kingdom; the Department of Cardiovascular Sciences (B.W.), School of Medicine, University of Leicester, Leicester, United Kingdom; and the First Department of Cardiology (C.V.), Athens Medical School, Hippokraton Hospital, Athens, Greece.

C.V. was writing coordinator for this article.

Correspondence to Charalambos Vlachopoulos, Kerassoundos 17, Athens 11528, Greece. E-mail cvlachop@otenet.gr

(*Hypertension*. 2007;50:154-160.)

© 2007 American Heart Association, Inc.

Hypertension is available at <http://www.hypertensionaha.org>

DOI: 10.1161/HYPERTENSIONAHA.107.090068




Figure 1. Change in contours in pressure wave (top) and flow wave (bottom) between the ascending aorta and the saphenous artery (reprinted from Reference 3).

in turn depends on the balance between vasoconstriction and vasodilatation in the peripheral circulation.

Another important consideration regarding the relationship between brachial and central aortic pressure is “pressure wave amplification.” Typically, the diastolic and mean pressures change little across the arterial tree. However, systolic BP is amplified when moving from the aorta to the periphery (Figure 1).³

Consequently, the pulsatile components of central and peripheral pressures (systolic BP and pulse pressure [PP]) may vary significantly. In general, brachial systolic and PP tend to overestimate central systolic and PP, especially in younger subjects who have more pronounced amplification. Substantial differences between central and brachial BP are also often also seen in older people, especially with tachycardia, exercise, use of vasoactive agents, or in those with systolic heart failure.

Techniques to Assess Central Hemodynamics

Although central pressures are ideally measured directly by using invasive devices, several methods have been devised currently to derive central pressures from analyses of applanated carotid and radial pulses or carotid distension waves.⁶ Among several commercial and noncommercial devices,^{3,6-11} the most widely used in clinical studies is the SphygmoCor device (AtCor Medical), which uses radial or carotid pulses and a validated generalized transfer function to estimate central pressures from the peripheral signal (Figure 2). When the applanated radial waveform is calibrated using direct intra-arterial pressures, the SphygmoCor calculations of aortic pressures are accurate,^{8,12} whereas the accuracy of this, as well as of other peripheral artery methods, decreases when the radial pulses are calibrated using inaccurate cuff pressures.¹³ This is also the case for carotid artery techniques. Alternatively, because mean BP remains virtually constant from central to peripheral arteries, the mean BP value computed from the area of the applanated carotid artery waveform can be calibrated using the mean BP obtained at the brachial artery level. Values of carotid pressures can then be derived.^{6,11} It should be emphasized that even when central




Figure 2. Central pressure waveform. The height of the late systolic peak above the inflection defines the augmented pressure, and the ratio of augmented pressure to PP defines the augmentation index (in percentage; modified from Reference 6).

pressures cannot be accurately calculated, these methods can display several features of the aortic (or carotid) pulse that are not dependent on the absolute values of BP, such as amplification of the pulse wave between central and peripheral artery, augmentation index, and arrival time of the reflected wave (the latter 2 are, however, dependent on identification of inflection point). Furthermore, because these features are also influenced by factors including heart rate, height, and age, modern computational capability could allow weighting of these factors in appropriate formulas. Potential problems related to the widespread application of arterial tonometry include appropriate training, supervision, and quality control, as they do for sphygmomanometry.

Arterial wave reflections indices and central pressures coupled with measurement of aortic pulse wave velocity, a direct measure of arterial stiffness, constitute a comprehensive and integrated approach of the study of arterial function.⁶ This is further highlighted by a possible differential ability to predict risk (augmentation index is better in younger persons and pulse wave velocity is better in older persons¹⁴) or by situations where, contrary to the usual behavior, pulse wave velocity and wave reflection indices move in opposite directions.¹⁵

Central Pressures and Central Indices as Markers and Predictors of Disease

Central hemodynamic variables have been shown to be independently associated with organ damage, incident cardiovascular disease, and events both in the general population and in various disease states.^{2,16-40} Tables 1 and 2 show such associations according to population studied, variable measured, and site and mode (invasive or noninvasive) of measurement.

Central Pressures and Central Indices as Markers of Disease and Predictors of Surrogate End Points

Increased aortic augmentation index is associated with coronary artery disease.²⁶ Central pressures also correlate with cardiovascular risk not only in patients with atherosclerotic disease but also in apparently healthy subjects.²² The late systolic augmentation of the central pressure waveform is associated with an increase in left ventricular mass index

TABLE 1. Cross-Sectional and Longitudinal Studies Indicating the Independent Value of Central Hemodynamics as Markers of Disease and Predictors of Surrogate Cardiovascular End Points

Source	Year, Country	Population	Design	Parameter	End Point
Saba et al ^{16*} †	1993, United States	Normotensives	Cross-sectional	Carotid Alx	LVMI, carotid thickness
Boutouyrie et al ^{17*} †	1999, France	Hypertensives	Cross-sectional	Carotid PP	Carotid thickness
Boutouyrie et al ^{18*} †	2000, France	Hypertensives	Longitudinal (9-month FU)	Carotid PP	Carotid IMT reduction with treatment
Roman et al ¹⁹	2000, United States	Normotensives, Hypertensives	Cross-sectional	Carotid systolic BP	Relative LV wall thickness
Waddell et al ^{20*} †	2001, Australia	CAD	Cross-sectional	Carotid BP	Extent of CAD
Nishijima et al ^{21*}	2001, Japan	Suspected CAD	Cross-sectional	Aortic fractional PP	Incident CAD
Nurnberger et al ²²	2002, Germany	Healthy+CVD	Cross-sectional	Carotid Alx	CV risk scores
Philippe et al ^{23*}	2002, France	CAD	Cross-sectional	Aortic PP	Extent of CAD
Hayashi et al ^{24*}	2002, Japan	Suspected CAD	Cross-sectional	Aortic Alx	Incident CAD
De Luca et al ²⁵ †	2004, REASON Study	Hypertensives	Longitudinal (1-year FU)	Carotid PP	LVMI reduction
Weber et al ²⁶	2004, Austria	Suspected CAD	Cross-sectional	Aortic AP, Alx	Incident CAD
Jankowski et al ^{27*}	2004, Poland	CAD	Cross-sectional	Aortic BP	Extent of CAD
Danchin et al ^{28*}	2004, France	Suspected CAD	Cross-sectional	Aortic PP	Incidence and extent of CAD
Booth et al ²⁹	2004, United Kingdom	Systemic vasculitis	Cross-sectional	Aortic Alx	Disease activity
Roman et al ³⁰	2007, United States	High-risk	Cross-sectional	Aortic PP	Carotid IMT and mass
Hashimoto et al ³¹ †	2007, Japan	Hypertensives	Longitudinal (1-year FU)	Aortic Alx	LVMI reduction with treatment

Alx indicates augmentation index; CAD, coronary artery disease; CV, cardiovascular; FU, follow-up; IMT, intima-media thickness; LV, left ventricular; LVMI, left ventricular mass index.

*Central pressure measured directly.

†These studies have shown incremental value of central indices over peripheral BP.

independent of age and mean BP,^{16,31} and carotid systolic BP is an independent determinant of left ventricular wall thickness.¹⁹ Moreover, central pressure is also more closely related to other important cardiovascular intermediate end points, such as vascular hypertrophy, extent of carotid atherosclerosis,^{17,30} and the ascending aorta diameter in patients with Marfan syndrome⁴¹ than brachial pressure. Recently, higher

carotid pressure augmentation in older individuals has been linked to increased flow pulsations entering the cerebral circulation, which may increase the risk of cerebral microvascular damage.⁴² In inflammatory disorders such as systemic vasculitis, augmentation index is a marker of disease activity, and it is independently associated with levels of C-reactive protein.²⁹

TABLE 2. Longitudinal Studies Indicating the Independent Value of Central Hemodynamics as Predictors of Events

Source	Year, country	Population	Design	Parameter	End Point
Nakayama et al ^{32*}	2000, Japan	CAD-PTCA	Longitudinal (3-mo FU)	Aortic fractional PP	Restenosis
Lu et al ^{33*}	2001, China	CAD-PTCA	Longitudinal (6-mo FU)	Aortic PP	Restenosis
London et al ³⁴ †	2001, France	ESRD	Longitudinal (52-mo FU)	Carotid Alx	CV mortality
Safar et al ² †	2002, France	ESRD	Longitudinal (52-mo FU)	Carotid PP, PP amplification	All-cause and CV mortality
Ueda et al ^{35*}	2004, Japan	CAD-PTCA	Longitudinal (6-mo FU)	Aortic Alx	Restenosis
Chirinos et al ^{36*} †	2005, United States	CAD	Longitudinal (3.2-y FU)	Aortic AP	CV mortality and events
Weber et al ³⁷ †	2005, Austria	CAD-PTCA	Longitudinal (2-y FU)	Aortic Alx	CV mortality and events
Dart et al ³⁸	2006, Australia	Elderly female hypertensives	Longitudinal (4.1-y FU)	Carotid Alx, brachial BP	CV mortality and events
Williams et al ³⁹ †	2006 CAFE Study	Hypertensives	Longitudinal (\leq 4-y FU)	Aortic PP	CV mortality and events during treatment
Roman et al ^{30,40} †	2005 and 2007, United States	High-risk	Longitudinal (4.8-y FU)	Aortic PP	CV mortality and events

ESRD indicates end-stage renal disease; PTCA, percutaneous transluminal coronary angioplasty; Alx, augmentation index; CAD, coronary artery disease; CV, cardiovascular; FU, follow-up.

*Central pressure measured directly.

†These studies have shown incremental value of central indices over peripheral BP.




Figure 3. Metaregression line relating the within-trial difference in systolic BP to the odds ratios for stroke. Odds ratios were calculated for experimental vs reference treatment. BP differences were obtained by subtracting achieved levels in experimental groups from those in reference groups. At left, pressure is the brachial measured, whereas at right, pressure is the estimated central pressure allowed for the decrease over brachial pressure of 3 mm Hg when the active therapy was a drug that reduces peripheral wave reflection (calcium channel blocker, angiotensin-converting enzyme inhibitor, or angiotensin receptor blocker). The curvilinear metaregression line (left) becomes linear (right) when central rather than peripheral pressures are plotted against odds ratios, indicating a direct relation between cardiovascular outcome and pressure changes induced by antihypertensive treatment (from References 3 and 45).

Central Pressures and Central Indices as Predictors of Events

Recent data from the Strong Heart Study confirm that peripheral PP, a simple index of arterial stiffness, is associated with a higher cardiovascular mortality independent of traditional risk factors, left ventricular hypertrophy, and reduced ejection fraction in adults without overt coronary heart disease.⁴³ Furthermore, data from the same study showed in a 5-year follow-up that the noninvasively determined central PP better predicts incident cardiovascular disease than does the corresponding brachial PP, possibly because of a more accurate representation of the vascular load on the left ventricle.^{30,40} The predictive value of central PP is significant even when subclinical atherosclerosis is taken into account.^{30,40} Central pressures and wave reflection indices are also strong independent predictors of all-cause and cardiovascular mortality in patients with end-stage renal failure.^{2,34} Moreover, in patients with coronary artery disease, wave reflections as expressed by central augmented pressure are powerful and independent predictors of recurrent acute coronary events or death.³⁶ Pulsatility of the ascending aortic pressure waveform is a powerful predictor of restenosis after angioplasty.^{32,33} Moreover, increased arterial wave reflections expressed by augmentation index are independently associated with an increased risk of severe short- and long-term cardiovascular events in patients undergoing percutaneous coronary interventions.³⁷ In contrast to the aforementioned evidence, a recent study conducted in elderly hypertensive women found that carotid augmentation index is not predictive of outcome.³⁸ However, the large Conduit Artery Function Evaluation (CAFE) Study reported that central PP derived from radial artery applanation tonometry independently predicts outcome in treated patients with hypertension.³⁹ Highlighting the interplay of small and large arteries in

risk determination, the structure of small arteries (which compose the main peripheral reflecting sites) was a predictor of events in patients with hypertension.⁴⁴

It should be noted, however, that evidence for a specific central pressure component or index does not necessarily apply to the others, and, thus, they should not be used interchangeably. Furthermore, there is clearly a need for additional studies of central pressures and indices in more general populations or in other disease states. Finally, it would be desirable for future studies to address whether central pressures provide incremental and independent prognostic value over other emerging biomarkers.

Implications for Therapy

BP reduction, per se, is the major determinant of the benefit of antihypertensive treatment. This has been shown by data from placebo-controlled trials, trials that compared more intensive versus less intensive BP-lowering strategies, and trials comparing different active regimens, and it is further supported by large meta-analyses of studies on antihypertensive treatment^{1,45-47} (Figure 3).

However, during the last decade, important multicenter trials gave rise to the hypothesis that new antihypertensive drugs, such as blockers of the renin-angiotensin system, may reduce cardiovascular outcomes beyond (peripheral) BP control. Particularly, in the Heart Outcomes Prevention Evaluation (HOPE), Losartan Intervention For Endpoint reduction in hypertension (LIFE), and the Australian National Blood Pressure 2 (ANBP2) studies, the observed clinical benefit tended to be greater than that expected from the decrease in peripheral BP. These potential effects “beyond BP control” are perhaps accounted for by protective properties of different drugs that affect subclinical organ damage or intermediate end points, such as arterial properties or central BP, for which

there is evidence that they are related to cardiovascular morbidity and mortality.^{1,48} Effects on central pressures may not be evident by pressure measurements in the periphery, because the reflected wave is added to a different part of the central waveform.⁴⁹ This may explain why drugs with similar reduction in peripheral pressures have a differential impact on cardiovascular outcomes. In this context, a short-term (4-week treatment) study showed that, in contrast to β -blocking drugs, angiotensin-converting enzyme inhibitors and calcium blockers may be more effective in reducing aortic systolic BP than that indicated by brachial pressure measurements.⁵⁰ Furthermore, more recent studies comparing the acute effects of angiotensin-converting enzyme inhibitors (ramipril) and the short-term (6-week) effects of angiotensin receptor blockers (eprosartan) with atenolol showed that, for a similar fall of brachial pressure, inhibitors of the renin-angiotensin axis may cause a significantly greater fall of central pressures.^{51,52} Interestingly, the pREterax in regression of Arterial Stiffness in a contrOLled double-bliNd (REASON) trial, which compared a perindopril-indapamide combination against atenolol, showed that normalization of brachial systolic BP is achieved with a significantly greater reduction of carotid systolic BP after a 12-month treatment with the combination.⁵³ In this study, compared with atenolol, the perindopril-indapamide combination was associated with a greater fall in left ventricular mass, and this was related to carotid but not brachial BP.²⁵

The notion that BP-lowering drugs can have substantially different effects on central aortic pressures and hemodynamics despite a similar impact on brachial BP and that central rather than peripheral BP should be a treatment target in trials was highlighted by the recent CAFE Study,³⁹ a major substudy within the Anglo-Scandinavian Cardiac Outcome Trial, which examined the impact of a conventional regimen (atenolol \pm thiazide) versus a contemporary one (amlodipine \pm perindopril). In the CAFE Study, despite similar effects in peripheral BP, there were substantial reductions in central aortic pressures and hemodynamic indices in favor of the amlodipine-based treatment. Because there were no differences in carotid-femoral pulse wave velocity and the pressure at the inflection point (height of P1, studied in a subgroup of patients), these changes were predominantly attributed to lower pressure wave augmentation because of decreased pulse wave reflection. Central aortic PP was a determinant of clinical outcome in this study.³⁹ These findings are in line with the enhancement of the relationship between events and BP reduction when allowance is made for a greater reduction in central than brachial pressure occurring with arterial dilating drugs (Figure 3, right). Because β -blockers are less effective than other drugs in preventing major cardiovascular events, especially stroke, the recent recommendations for the treatment of hypertension suggest against their use for initial treatment of hypertension.^{54,55} It is currently unknown whether this is a class effect (including those β -blockers with vasodilating properties) or whether this only pertains to atenolol, the main β -blocker used in clinical trials of hypertension.

Other studies have also shown that surrogate end points can be predicted by indices of central hemodynamics after

therapy. Aortic and radial augmentation index, as well as amplification of the pulse between central and radial arteries, were superior to conventional cuff measures in predicting reduction in left ventricular mass during antihypertensive therapy.³¹ Furthermore, the reduction in carotid wall diameter and hypertrophy with antihypertensive treatment was related to carotid PP but not to mean BP.¹⁸

Implementation in Clinical Practice

Both peripheral and central BP fulfil the criteria of a surrogate marker (type 2 marker).⁵⁶ Accordingly, regarding the theoretical basis, central BP is pathophysiologically more relevant than the corresponding peripheral BPs. Furthermore, current, noninvasive, validated, and easy-to-use techniques can estimate central BPs and wave reflection indices. Reproducibility is very good (please see <http://hyper.ahajournals.org>), but improvement is desirable. In addition, because peripheral BP is perhaps the most firmly established cardiovascular risk predictor, the question of whether central BP provides incremental value over and above peripheral BP is pivotal. Evidence is growing that this may be indeed the case, but more data are needed with regard to each central pressure index.^{2,30,34,36,40,43,55} Finally, the CAFE Study³⁹ has provided the first randomized, prospective event-based evidence that central BP and related indices may be a useful guide to treatment. Nevertheless, more studies are needed.

Before implementation into clinical practice, determination of normal values for wave reflections indices and central BPs is mandatory. Importantly, the recently published Anglo-Cardiff Collaborative Trial¹⁴ and the European Project on Genes in Hypertension, as well as ongoing studies (European Network for Noninvasive Determination of Large Arteries, National Institute of Aging, and Framingham) are making important progress in this direction.

Perspectives

The prognostic value of brachial BP is well established. However, there is mounting evidence suggesting that central pressure and indices correlate more closely with intermediate measures of cardiovascular risk than brachial pressure and that they independently predict future cardiovascular events. Moreover, despite similar effects on brachial pressure, antihypertensive drugs have differential effects on central pressure. This may explain the superiority of vasodilating drugs in recent outcome trials. However, extension of the existing data regarding the superiority of central pressures over and above brachial BP in a wider range of populations and disease states is desirable. Inclusion of a parameter in patient assessment and management should serve various purposes, such as advancement of science, physician education, and practicality of use in a range of settings, whereas cost should also be taken into consideration. Assessment of central pressures meets these criteria to a varying degree at present. *Definition of terms such as central and peripheral BP, arterial stiffness, wave reflections, and systolic BP and PP amplification should be readily available to both clinicians and researchers and introduced in the guidelines on hypertension and cardiovascular risk.* Although brachial BP remains our point of reference, there is a definite sense that efforts to investigate

and advance from the status quo of cuff BP measurements should be pursued. Acquired evidence of this sort would allow for supportable recommendations to the committees whose responsibility it is to issue guidelines for the evaluation and management of hypertension.

Disclosures

M.F.O. is the founding director of AtCor Medical, the manufacturer of systems for analyzing the arterial pulse. The remaining authors report no conflicts.

References

- Guidelines Committee. 2003 European Society of Hypertension-European Society of Cardiology guidelines for the management of arterial hypertension. *J Hypertens*. 2003;21:1011–1053.
- Safar ME, Blacher J, Pannier B, Guerin AP, Marchais SJ, Guyonvarc'h PM, London GM. Central pulse pressure and mortality in end-stage renal disease. *Hypertension*. 2002;39:735–738.
- Nichols WW, O'Rourke MF. *McDonald's Blood Flow in Arteries*. London, United Kingdom: Arnold; 2005.
- Safar ME, O'Rourke MF. *Handbook of Hypertension: Arterial Stiffness and Wave Reflection*. Paris, France: Elsevier; 2005.
- Avolio AP, Chen SG, Wang RP, Zhang CL, Li MF, O'Rourke MF. Effects of aging on changing arterial compliance and left ventricular load in a northern Chinese urban community. *Circulation*. 1983;68:50–58.
- Laurent S, Cockcroft J, Van Bortel L, Boutouyrie P, Giannattasio C, Hayoz D, Pannier B, Vlachopoulos C, Wilkinson I, Struijker-Boudier H; European Network for Non-invasive Investigation of Large Arteries. Expert consensus document on arterial stiffness: methodological issues and clinical applications. *Eur Heart J*. 2006;27:2588–2605.
- O'Rourke MF, Adji A. An updated clinical primer on large artery mechanics: implications of pulse waveform analysis and arterial tonometry. *Curr Opin Cardiol*. 2005;20:275–281.
- Paucal AL, O'Rourke MF, Kon ND. Prospective evaluation of a method for estimating ascending aortic pressure from the radial artery pressure waveform. *Hypertension*. 2001;38:932–937.
- Mitchell GF, Parise H, Benjamin EJ, Larson MG, Keyes MJ, Vita JA, Vasan RS, Levy D. Changes in arterial stiffness and wave reflection with advancing age in healthy men and women. The Framingham Heart Study. *Hypertension*. 2004;43:1239–1245.
- Van Bortel LM, Balkestein EJ, van der Heijden-Spek JJ, Vanmolkot FH, Staessen JA, Kragten JA, Vredeveld JW, Safar ME, Struijker Boudier HA, Hoeks AP. Non-invasive assessment of local arterial pulse pressure: comparison of applanation tonometry and echo-tracking. *J Hypertens*. 2001;19:1037–1044.
- Verbeke F, Segers P, Heireman S, Vanholder R, Verdonck P, Van Bortel LM. Noninvasive assessment of local pulse pressure: importance of brachial-to-radial pressure amplification. *Hypertension*. 2005;46:244–248.
- Sharman JE, Lim R, Qasem AM, Coombes JS, Burgess MI, Franco J, Garrahy P, Wilkinson IB, Marwick TH. Validation of a generalized transfer function to noninvasively derive central blood pressure during exercise. *Hypertension*. 2006;47:1203–1208.
- Smulyan H, Siddiqui DS, Carlson RJ, London GM, Safar ME. Clinical utility of aortic pulses and pressures calculated from applanated radial-artery pulses. *Hypertension*. 2003;42:150–155.
- McEniery CM, Yasmin, Hall IR, Qasem A, Wilkinson IB, Cockcroft JR; ACCT Investigators. Normal vascular aging: differential effects on wave reflection and aortic pulse wave velocity: the Anglo-Cardiff Collaborative Trial (ACCT). *J Am Coll Cardiol*. 2005;46:1753–1760.
- Vlachopoulos C, Dima I, Aznaouridis K, Vasiliadou C, Ioakeimidis N, Aggeli C, Toutouza M, Stefanadis C. Acute systemic inflammation increases arterial stiffness and decreases wave reflections in healthy individuals. *Circulation*. 2005;112:2193–2200.
- Saba PS, Roman MJ, Pini R, Spitzer M, Ganau A, Devereux RB. Relation of arterial pressure waveform to left ventricular and carotid anatomy in normotensive subjects. *J Am Coll Cardiol*. 1993;22:1873–1880.
- Boutouyrie P, Bussy C, Lacolley P, Girerd X, Laloux B, Laurent S. Association between local pulse pressure, mean blood pressure, and large-artery remodeling. *Circulation*. 1999;100:1387–1393.
- Boutouyrie P, Bussy C, Hayoz D, Hengstler J, Dartois N, Laloux B, Brunner H, Laurent S. Local pulse pressure and regression of arterial wall hypertrophy during long-term antihypertensive treatment. *Circulation*. 2000;101:2601–2606.
- Roman MJ, Ganau A, Saba PS, Pini R, Pickering T, Devereux R. Impact of arterial stiffening on left ventricular structure. *Hypertension*. 2000;36:489–494.
- Waddell TK, Dart AM, Medley TL, Cameron JD, Kingwell BA. Carotid pressure is a better predictor of coronary artery disease than brachial pressure. *Hypertension*. 2001;38:927–931.
- Nishijima T, Nakayama Y, Tsumura K, Yamashita N, Yoshimaru K, Ueda H, Hayashi T, Yoshikawa J. Pulsatility of ascending aortic blood pressure waveform is associated with an increased risk of coronary heart disease. *Am J Hypertens*. 2001;14:469–473.
- Nurnberger J, Keflioglu-Scheiber A, Opazo Saez AM, Wenzel RR, Philipp T, Schafers RF. Augmentation index is associated with cardiovascular risk. *J Hypertens*. 2002;20:2407–2414.
- Philippe F, Chemaly E, Blacher J, Mourad JJ, Dibie A, Larrazet F, Laborde F, Safar ME. Aortic pulse pressure and extent of coronary artery disease in percutaneous transluminal coronary angioplasty candidates. *Am J Hypertens*. 2002;15:672–677.
- Hayashi T, Nakayama Y, Tsumura K, Yoshimaru K, Ueda H. Reflection in the arterial system and the risk of coronary heart disease. *Am J Hypertens*. 2002;15:405–409.
- de Luca N, Asmar RG, London GM, O'Rourke MF, Safar ME; REASON Project Investigators. Selective reduction of cardiac mass and central blood pressure on low-dose combination perindopril/indapamide in hypertensive subjects. *J Hypertens*. 2004;22:1623–1630.
- Weber T, Auer J, O'Rourke M, Kvas E, Lassnig E, Berent R, Eber B. Arterial stiffness, wave reflections, and the risk of coronary artery disease. *Circulation*. 2004;109:184–189.
- Jankowski J, Kawecka-Jaszcz K, Bryniarski L, Czarnecka D, Brzozowska-Kiszka M, Posnik-Urbanska A, Kopec G, Dragan J, Klecha A, Dudek D. Fractional diastolic and systolic pressure in the ascending aorta are related to the extent of coronary artery disease. *Am J Hypertens*. 2004;17:641–646.
- Danchin N, Benetos A, Lopez-Sublet M, Demicheli T, Safar M, Mourad JJ; ESCAPP Investigators. Aortic pulse pressure is related to the presence and extent of coronary artery disease in men undergoing diagnostic coronary angiography: a multicenter study. *Am J Hypertens*. 2004;17:129–133.
- Booth AD, Wallace S, McEniery CM, Yasmin, Brown J, Jayne DR, Wilkinson IB. Inflammation and arterial stiffness in systemic vasculitis: a model of vascular inflammation. *Arthritis Rheum*. 2004;50:581–588.
- Roman MJ, Devereux RB, Kizer JR, Lee ET, Galloway JM, Ali T, Umans JG, Howard BV. Central pressure more strongly relates to vascular disease and outcome than does brachial pressure: the Strong Heart Study. *Hypertension*. 2007;50:197–203.
- Hashimoto J, Imai Y, O'Rourke MF. Indices of pulse wave analysis are better predictors of left ventricular mass reduction than cuff pressure. *Am J Hypertens*. 2007;20:378–384.
- Nakayama Y, Tsumura K, Yamashita N, Yoshimaru K, Hayashi T. Pulsatility of ascending aortic pressure waveform is a powerful predictor of restenosis after percutaneous transluminal coronary angioplasty. *Circulation*. 2000;101:470–472.
- Lu TM, Hsu NW, Chen YH, Lee WS, Wu CC, Ding YA, Chang MS, Lin SJ. Pulsatility of ascending aorta and restenosis after coronary angioplasty in patients >60 years of age with stable angina pectoris. *Am J Cardiol*. 2001;88:964–968.
- London GM, Blacher J, Pannier B, Guerin A, Marchais S, Safar M. Arterial wave reflections and survival in end-stage renal failure. *Hypertension*. 2001;38:434–438.
- Ueda H, Hayashi T, Tsumura K, Yoshimaru K, Nakayama Y, Yoshikawa J. The timing of the reflected wave in the ascending aortic pressure predicts restenosis after coronary stent placement. *Hypertens Res*. 2004;27:535–540.
- Chirinos JA, Zambrano JP, Chakko S, Veerani A, Schob A, Willens HJ, Perez G, Mendez AJ. Aortic pressure augmentation predicts adverse cardiovascular events in patients with established coronary artery disease. *Hypertension*. 2005;45:980–985.
- Weber T, Auer J, O'Rourke MF, Kvas E, Lassnig E, Lamm G, Stark N, Rammer M, Eber B. Increased arterial wave reflections predict severe cardiovascular events in patients undergoing percutaneous coronary interventions. *Eur Heart J*. 2005;26:2657–2663.
- Dart AM, Gatzka CD, Kingwell BA, Willson K, Cameron JD, Liang YL, Berry KL, Wing LM, Reid CM, Ryan P, Beilin LJ, Jennings GL, Johnston CI, McNeil JJ, Macdonald GJ, Morgan TO, West MJ. Brachial blood

- pressure but not carotid arterial waveforms predict cardiovascular events in elderly female hypertensives. *Hypertension*. 2006;47:785–790.
39. Williams B, Lacy PS, Thom SM, Cruickshank K, Stanton A, Collier D, Hughes AD, Thurston H, O'Rourke M; CAFE Investigators; Anglo-Scandinavian Cardiac Outcomes Trial Investigators; CAFE Steering Committee and Writing Committee. Differential impact of blood pressure-lowering drugs on central aortic pressure and clinical outcomes. *Circulation*. 2006;113:1213–1225.
 40. Roman MJ, Kizer JR, Ali T, Lee ET, Galloway JM, Fabsitz RR, Henderson JA, Howard BV. Central blood pressure better predicts cardiovascular events than does peripheral blood pressure: the Strong Heart Study. Presented at the AHA Scientific Sessions, November 13–16, 2005. Abstract.
 41. Jondeau G, Boutouyrie P, Lacolley P, Laloux B, Dubourg O, Bourdarias JP, Laurent S. Central pulse pressure is a major determinant of ascending aorta dilation in Marfan syndrome. *Circulation*. 1999;99:2677–2681.
 42. Hirata K, Yaginuma T, O'Rourke MF, Kawakami M. Age-related change in the carotid artery flow and pressure pulses: implication to cerebral microvascular disease. *Stroke*. 2006;37:2552–2556.
 43. Palmieri V, Devereux RB, Hollywood J, Bella JN, Liu JE, Lee ET, Best LG, Howard BV, Roman MJ. Association of pulse pressure with cardiovascular outcome is independent of left ventricular hypertrophy and systolic dysfunction: The Strong Heart Study. *Am J Hypertens*. 2006;19:601–607.
 44. Rizzoni D, Porteri E, Boari GE, De Ciuceis C, Sleiman I, Muiesan ML, Castellano M, Miclini M, Agabiti-Rosei E. Prognostic significance of small-artery structure in hypertension. *Circulation*. 2003;108:2230–2235.
 45. Staessen JA, Wang JG, Thijs L. Cardiovascular protection and blood pressure reduction: a meta-analysis. *Lancet*. 2001;358:1305–1315.
 46. Wang JG, Li Y, Franklin SS, Safar M. Prevention of stroke and myocardial infarction by amlodipine and angiotensin receptor blockers: a quantitative overview. *Hypertension*. 2007;50:219–227.
 47. Blood Pressure Lowering Treatment Trialists' Collaboration. Effects of different blood pressure-lowering regimens on major cardiovascular events in individuals with and without diabetes mellitus: results of prospectively designed overviews of randomized trials. *Arch Intern Med*. 2005;165:1410–1419.
 48. Mancia G. Role of outcome trials in providing information on antihypertensive treatment: importance and limitations. *Am J Hypertens*. 2006;19:1–7.
 49. Vlachopoulos C, Hirata K, O'Rourke MF. Pressure-altering agents affect central aortic pressures more than is apparent from upper limb measurements in hypertensive patients: the role of arterial wave reflections. *Hypertension*. 2001;38:1456–1460.
 50. Morgan T, Lauri J, Bertram D, Anderson A. Effect of different antihypertensive drug classes on central aortic pressure. *Am J Hypertens*. 2004;17:118–123.
 51. Hirata K, Vlachopoulos C, Adji A, O'Rourke MF. Benefits from angiotensin-converting enzyme inhibitor 'beyond blood pressure lowering': beyond blood pressure or beyond the brachial artery? *J Hypertens*. 2005;23:551–556.
 52. Dhakam Z, McEniery CM, Yasmin, Cockcroft JR, Brown MJ, Wilkinson IB. Atenolol and eprosartan: differential effects on central blood pressure and aortic pulse wave velocity. *Am J Hypertens*. 2006;19:214–219.
 53. London GM, Asmar RG, O'Rourke MF, Safar ME, on behalf of the REASON Project Investigators. Mechanism(s) of selective systolic blood pressure reduction after a low-dose combination of perindopril/indapamide in hypertensive subjects: comparison with atenolol. *J Am Coll Cardiol*. 2004;43:92–99.
 54. The National Collaborating Centre for Chronic Conditions. Management of hypertension in adults in primary care: partial update. Royal College of Physicians Web site. Available at: <http://www.nice.org.uk/guidance/CG34>. Accessed December 28, 2006.
 55. Williams B. Evolution of hypertensive disease: a revolution in guidelines. *Lancet*. 2006;368:6–8.
 56. Vasan RS. Biomarkers of cardiovascular disease: molecular basis and practical considerations. *Circulation*. 2006;113:2335–2362.

Table: Reproducibility/repeatability and accuracy/validity of estimated central hemodynamic indices.

First author	Year, country	Population	Variables - reproducibility
REPRODUCIBILITY / REPEATABILITY			
Wilkinson ¹	1998, UK	Healthy, hypertensives, diabetics	Aortic AIx: intraobserver difference 0.49±5.37% interobserver difference 0.23±3.80%
Liang ²	1998, Australia	Healthy	Carotid AIx: coefficient of variation 1.3%
Siebenhoffer ³	1999, UK	Healthy	Derived aortic SBP: interobserver difference 0.1±1.7 mmHg Derived aortic DBP: interobserver difference 0.1±0.7 mmHg Aortic AIx: interobserver difference 0.4±6.4%
Filipovsky ⁴	2000, Czech Republic	Healthy	Aortic AIx: intraobserver variability 7.7% Aortic SBP: intraobserver variability 6.3%
Savage ⁵	2002, UK	Chronic renal failure	Aortic AIx: intraobserver difference 0±4% interobserver difference 0±3% Aortic MAP: intraobserver difference 0±3 mmHg interobserver difference 1±9 mmHg
Matsui ⁶	2004, Japan	Hypertensives	Carotid AIx: intraobserver difference 0.5±5.9%
Papaioannou ⁷	2004, Greece	Cardiogenic shock	Aortic AIx: intraobserver difference 0.10±5.82%
Williams (CAFÉ) ⁸	2006, UK-Sweden	Hypertensives	Aortic SBP: interobserver difference 0.3±2.9 mmHg Aortic AIx: interobserver difference 1.5±5.9%
Roman (Strong Hear Study) ^{9,10}	2007, USA	High-risk subjects	Carotid SBP: interobserver CC: 0.99 intraobserver CC: 0.99 Carotid DBP: interobserver CC: 0.99 intraobserver CC: 0.99
Paucá (radial) ¹¹	2001, Australia	Cardiac surgery (mostly CABG)	Resting conditions: SBP 0.0±4.4 mmHg; DBP 0.6±1.7 mmHg; PP-0.7±4.2 mmHg; MAP -0.5±2.0 mmHg Nitroglycerin infusion: SBP -0.2±4.3 mmHg; DBP 0.6±1.7 mmHg; PP -0.8±4.1 mmHg; MAP -0.4±1.8 mmHg.

Table: (continued)

First author	Year, country	Population	Variables - reproducibility
ACCURACY / VALIDITY			
(differences between calculated central BP with tonometry and measured central BP with catheterization)			
Van Bortel (carotid) ¹²	2001, Belgium, Netherlands	Coronary angiography	PP -10.2±14.3 mmHg
Smulyan (radial) ¹³	2003, USA- France	Coronary angiography	SBP 1.5±11.3 mmHg; DBP -10.4±12.7 mmHg; PP 11.5±13.6 mmHg
Hope (radial) ¹⁴	2004, Australia	Cardiac catheterization	SBP (invasive calibration) -1±8 mmHg SBP (non-invasive calibration) 7±12 mmHg
Sharman (radial) ¹⁵	2006, Australia	Coronary angiography	Rest SBP -1.3±3.2 mmHg Exercise SBP -4.7±3.3 mmHg

AIx indicates augmentation index; CC, correlation coefficient; DBP, diastolic blood pressure; MAP, mean arterial pressure; PP, pulse pressure; SBP, systolic blood pressure.

Reproducibility / repeatability of measurements with a single method expresses the error in measurements that is accounted for by biological variation of the variable under study or by the conditions of measurement (environment and technical skills).

Accuracy / validity of a method expresses the error of measurements with the method under study compared with another method that is considered a “gold standard”.

REFERENCES.

1. Wilkinson IB, Fuchs SA, Jansen IM, Spratt JC, Murray GD, Cockcroft JR, Webb DJ. Reproducibility of pulse wave velocity and augmentation index measured by pulse wave analysis. *J Hypertens*. 1998;16:2079–2084.
2. Liang YL, Teede H, Kotsopoulos D, Shiel L, Cameron JD, Dart AM, McGrath BP. Non-invasive measurements of arterial structure and function: repeatability, interrelationships and trial sample size. *Clin Sci*. 1998;95:669–679.
3. Siebenhofer A, Kemp C, Sutton A, Williams B. The reproducibility of central aortic blood pressure measurements in healthy subjects using applanation tonometry and sphygmocardiography. *J Hum Hypertens*. 1999;13:625-629.
4. Filipovsky J, Svobodova V, Pecen L. Reproducibility of radial pulse wave analysis in healthy subjects. *J Hypertens* 2000; 18:1033-1040.
5. Savage MT, Ferro CJ, Pinder SJ, Tomsom CR. Reproducibility of derived central arterial waveforms in patients with chronic renal failure. *Clin Sci*. 2002; 103:59–65.
6. Matsui Y, Kario K, Ishikawa J, Eguchi K, Hoshide S, Shimada K. Reproducibility of arterial stiffness indices (pulse wave velocity and augmentation index) simultaneously assessed by automated pulse wave analysis and their associated risk factors in essential hypertensive patients. *Hypertens Res*. 2004;27:851–857.
7. Papaioannou TG, Stamatelopoulos KS, Gialafos E, Vlachopoulos C, Karatzis E, Nanas J, Lekakis J. Monitoring of arterial stiffness indices by applanation tonometry and pulse wave analysis: reproducibility at low blood pressures. *J Clin Monit*. 2004;18:137–144.
8. Williams B, Lacy PS, Thom SM, Cruickshank K, Stanton A, Collier D, Hughes AD, Thurston H, O'Rourke M; CAFE Investigators; Anglo-Scandinavian Cardiac Outcomes Trial Investigators; CAFE Steering Committee and Writing Committee. Differential impact of blood pressure-lowering drugs on central aortic pressure and clinical outcomes. *Circulation*. 2006;113:1213-1225.
9. Roman MJ, Saba PS, Pini R, Spitzer M, Pickering TG, Rosen S, Alderman MH, Devereux RB. Parallel cardiac and vascular adaptation in hypertension. *Circulation*. 1992; 86:1909-1918.

10. Roman MJ, Devereux RB, Kizer JR, Lee ET, Galloway JM, Ali T, Fabsitz RR, Henderson JA, Howard BV. Central aortic pressure better predicts outcome and better correlates with vascular disease than does brachial pressure: The Strong Heart Study. Sixth International Workshop on “Structure and Function of the Vascular System”, Paris, February 1-3, 2007 [abstract].
11. Pauca AL, O’Rourke MF, Kon ND. Prospective evaluation of a method for estimating ascending aortic pressure from the radial artery pressure waveform. *Hypertension*. 2001;38:932-937.
12. Van Bortel LM, Balkestein EJ, van der Heijden-Spek JJ, Vanmolkot FH, Staessen JA, Kragten JA, Vredeveld JW, Safar ME, Struijker Boudier HA, Hoeks AP. Non-invasive assessment of local arterial pulse pressure: comparison of applanation tonometry and echo-tracking. *J Hypertens*. 2001;19:1037-1044.
13. Smulyan H, Siddiqui DS, Carlson RJ, London GM, Safar ME. Clinical utility of aortic pulses and pressures calculated from applanated radial-artery pulses. *Hypertension*. 2003;42:150-155.
14. Hope SA, Meredith IT, Cameron JD. Effect of non-invasive calibration of radial waveforms on error in transfer-function-derived central aortic waveform characteristics. *Clin Sci*. 2004;107:205–211.
15. Sharman JE, Lim R, Qasem AM, Coombes JS, Burgess MI, Franco J, Garrahy P, Wilkinson IB, Marwick TH. Validation of a generalized transfer function to noninvasively derive central blood pressure during exercise. *Hypertension*. 2006;47:1203-1208.